

Orange-Senqu River Awareness Kit

The River Basin

- Introduction
- ▶ Geography
- ▶ Climate and Weather
- ▶ Hydrology
- ▶ Water Quality
- ▶ **Ecology and Biodiversity**
- Ecology
- Aquatic Ecology
- Building Blocks
- Aquatic Habitats
- Life in Aquatic Ecosystems
- Food Chains and Webs
- **Biomass and Production**
- Classification of Organisms
- Factors Affecting Ecosystems
- Wetlands
- Biodiversity
- Millennium Ecosystem Assessment
- References



Feedback

[send a general website comment](#)

[report a specific problem with this page](#)

Ecology and Biodiversity: Life in Aquatic Ecosystems: Biomass and Production

Organisms use energy to maintain biological functions and to enable growth and reproduction. The organic matter produced by autotrophs and heterotrophs, in excess of what they need to sustain life, adds to the ecosystem's total biomass. The biomass in an ecosystem includes the mass of all living and dead organic matter. Production is the incremental increase in biomass produced by organisms over a period of time. Estimates of biomass and production can be used to assess the health of aquatic ecosystems.


Primary production refers to the production of organic matter, such as new cells, mainly by photosynthetic plants. It is expressed as a rate of biomass production—for example, the amount of wood produced each year.

Secondary production is the assimilation of organic material and building of tissue by heterotrophs, and may involve animals eating plants, animals eating other animals, or microorganisms decomposing dead organisms to obtain the resources (material, energy, nutrients) needed to produce biomass. Secondary production is also expressed as a rate of biomass production, such as the amount of meat produced by grazing cattle each year.

In a productive environment, living plant or animal tissue will accumulate over time. Biomass is the amount of this accumulated material at a given moment, while production is the rate of increase in the total biomass. In a river system, biomass may be lost by export (such as downstream transport), or gained by import from other systems (such as leaves falling into a stream).

Estimates of biomass and production can be applied at various spatial scales and to broad or narrow groups of organisms, such as all organisms in a lake, all fish in a lake, or all yellow fish in a lake. Production is often difficult to estimate, since it requires, among other things, accurate measurements of biomass, repeated at consistent intervals over a long period.

High biomass does not necessarily imply high production, and vice versa. For example, the biomass of plankton in a waterbody may be low, but because plankton grow and reproduce quickly, the plankton population may replace itself relatively quickly—it has a high rate of production.



This figure illustrates energy flows related to the accumulation of biomass in an individual animal. Energy is obtained through food, while energy is consumed or lost in metabolic processes (i.e., respiration) and the excretion of wastes.

Source: NTEAP 2007
(click to enlarge)

[Next: Classification of Organisms](#)

Interactive

Basin Map

Explore the sub-basins of the Orange-Senqu River

[enter](#)

Video Tour

Tour video scenes along the Orange-Senqu River related to the River Basin

[enter](#)

Geography Maps

Investigate land cover and terrestrial ecoregions in the basin

[enter](#)

Water Cycle

Examine how the hydrologic cycle moves water through and around the earth

[enter](#)

Food Web

Explore the interactions of living organisms in aquatic environments

[enter](#)