


Orange-Senqu River Awareness Kit

[HOME](#)
[THE RIVER BASIN](#)
[PEOPLE AND THE RIVER](#)
[GOVERNANCE](#)
[RESOURCE MANAGEMENT](#)

The River Basin →

- Introduction
- ▶ Geography
- ▶ Climate and Weather
- ▶ Hydrology
- ▶ Water Quality
- ▼ **Ecology and Biodiversity**
- Ecology
- Aquatic Ecology
- Building Blocks
- Aquatic Habitats
- **Life in Aquatic Ecosystems**
- Food Chains and Webs
- Biomass and Production
- Classification of Organisms
- Microorganisms
- Plants
- Invertebrates
- Vertebrates
- Fish Species
- Factors Affecting Ecosystems
- Wetlands
- Biodiversity
- Millennium Ecosystem Assessment
- References



Feedback

[send a general website comment](#)
[report a specific problem with this page](#)

Ecology and Biodiversity: Aquatic Ecology: **Life in Aquatic Ecosystems**

The types of organisms inhabiting aquatic ecosystems depends on many factors including available resources and the physical environment. Biological communities—the species present and their relative abundance—arise through interactions between different types of organisms. These interactions can be partially described in terms of:

- [Food Chains and Food Webs](#)
- [Biomass and Production](#)

Basic Concepts - Energy and Food

In aquatic ecology, biologists often classify organisms according to how they obtain energy to live, grow and reproduce. Because sunlight is the ultimate source of energy for all organisms, a basic distinction lies between those who use its energy directly—**autotrophs**—and those who receive it indirectly by consuming other organisms—**heterotrophs**.

Autotrophs

Autotrophs, or **producers**, are organisms that can manufacture their own organic material from inorganic sources. Most autotrophs carry out this process using **photosynthesis**, the process by which plants and algae use solar energy to combine carbon dioxide with water to produce starch, sugars and oxygen.

Photosynthesis is the most important biological process on the planet, and its products drive the biological activity of nearly all ecosystems, including aquatic environments. The oxygen produced is available to be used by other organisms, making photosynthesis an important controller of carbon dioxide and oxygen in any environment.

Photosynthesis in aquatic systems is carried out by a wide variety of autotrophs, which range in size from microscopic single-celled organisms to large aquatic plants (macrophytes). Autotrophs are primary producers, because they produce the first level of organic compounds from simple inorganic substances. Ultimately, all other types of organisms (heterotrophs) are dependent on the organic carbon produced by autotrophs.

Because photosynthesis depends on sunlight, the distribution of autotrophs depends in part on the amount of light available in an aquatic ecosystem. In shallow, stony rivers, periphyton (or biofilm) – especially diatoms and cyanobacteria – are the main primary producers, but shade from riparian vegetation can limit photosynthesis; nutrients may also be in short supply in these habitats. In wider rivers, reduced shading from riparian vegetation allows the river surface to receive more light. However, in deep or turbid sections, light penetration may be insufficient to sustain growth of autotrophs.

Heterotrophs

Heterotrophs, or **consumers**, are organisms that must obtain energy by consuming other organisms (autotrophs or other heterotrophs) as food. From the perspective of energy flow in ecological systems, heterotrophs can be classified according to what they eat:

- **Herbivores** are called **primary consumers** because they eat only plants
- **Carnivores** are called **secondary consumers** because they feed on other animals
- **Omnivores** feed both on autotrophs and other heterotrophs; that is, they eat both plants and animals. Many aquatic organisms, including fish, are omnivorous
- **Detritivores** consume dead organic matter (detritus). Detritivores include many bacteria and fungi, invertebrates such as worms and insects, and some scavenging vertebrates. Aquatic insects, for instance, shred dead leaves, but also consume bacteria and fungi growing on the leaves

Heterotrophs can also be classified according to how they obtain food energy (i.e., functional feeding groups), and by their specific roles in the aquatic ecosystem (Cummins and Klug 1979):

- **Grazer-scrapers** feed on periphyton and biofilm
- **Shredders** are detritivores feeding on coarse organic particles, especially leaf litter derived from the riparian zone
- **Collectors** eat fine organic particles and can be subdivided according to whether the food particles they collect are *suspended* in the water (e.g., filtering-collectors or filter-feeders), or have been *deposited on the substratum* (collector-gatherers)
- **Deposit-feeders** ingest fine bottom sediments and the organic material that they contain
- **Predators** eat other animals

I Interactive

Basin Map

Explore the sub-basins of the Orange-Senqu River

[enter](#)

Video Tour

Tour video scenes along the Orange-Senqu River related to the River Basin

[enter](#)

Geography Maps

Investigate land cover and terrestrial ecoregions in the basin

[enter](#)

Water Cycle

Examine how the hydrologic cycle moves water through and around the earth

[enter](#)

Food Web

Explore the interactions of living organisms in aquatic environments

[enter](#)



All birds, including this Cape Cormorant, are heterotrophs because they cannot harness the sun's energy.

Source: MBoY68 2007
(click to enlarge)

[Next: Food Chains and Food Webs](#) ►

You are here: [Home Page](#) > [The River Basin](#) > [Ecology and Biodiversity](#) > [Aquatic Ecology](#) > [Life in Aquatic Ecosystems](#)